

**COUNTY OF SACRAMENTO
EMERGENCY MEDICAL SERVICES AGENCY**



Program Document: **Traumatic Cardiac Arrest**

Policy Number: 8032.02

Initial Date: 06/22/21

Last Approved Date: 09/14/23

Effective Date: 05/01/24

Next Review Date: 03/01/25

Signature on File

EMS Medical Director

Signature on File

EMS Administrator

Purpose:

- A. To serve as the treatment standard for treating traumatic cardiac arrest patients.

Authority:

- A. California Health and Safety Code, Division 2.5
- B. California Code of Regulations, Title 22, Division 9

Protocol:

- A. The pathophysiology of traumatic cardiac arrest differs from medical cardiac arrest and is primarily due to one or a combination of factors: hypovolemia, obstruction of blood flow, and hypoxia.
- B. The initial cardiac rhythm for most patients in survivable traumatic cardiac arrest is pulseless electrical activity (PEA). Traumatic cardiac arrest PEA is most often a very low output state due to hypovolemia.
- C. Traumatic cardiac arrest patients undergoing resuscitation shall be transported as quickly as possible to the hospital.
- D. Patients with trauma in cardiac arrest who, by prehospital presentation, may have suffered a medical event before trauma shall undergo medical cardiac arrest resuscitation per Policy# 8031 - Cardiac Arrest, with attention and appropriate management to emergent trauma needs (hemorrhage control, pneumothorax decompression as indicated, and orthopedic immobilization as indicated)
- E. There is no evidence-based medical support for the use of medications in traumatic cardiac arrest. In traumatic arrest, Epinephrine and Amiodarone are NOT indicated in traumatic cardiac arrest. If there is any doubt as to the cause of the arrest, treat it as a non-traumatic arrest.

Policy:**BLS:**

1. Treat immediate threats to life
2. External hemorrhage control per PD# 8065 - Hemorrhage Control. Apply tourniquets as necessary.
3. Airway and Breathing: Clear airway when indicated; place OPA and BVM ventilation.
4. Chest compressions/high-quality CPR for any rhythm other than Wide Complex PEA < 40 bpm or Asystole.
5. The use of a Mechanical CPR Device should be omitted if it will cause a delay in transport.
6. Expedite transport to the closest Trauma Center.

ALS:

1. Continue transport with BLS airway if adequate ventilation/chest rise is achieved. Advanced airway as needed per policy.
2. Correct potential obstructive shock – maintain a high index of suspicion for tension pneumothorax. Bilateral needle thoracostomy per PD# 8015 – Trauma.
3. Obtain large-bore IV or IO access. Give 1 liter of Normal Saline bolus by pressure bag infusion.
4. Cardiac monitoring – defibrillate shockable rhythms.

Post Resuscitation Consideration:

- A. Any traumatic cardiac arrest patient who has a Return of Spontaneous Circulation (ROSC) during any part of the resuscitation and who is transported shall be transported to a Trauma Center.
- B. Intravenous (IV) or Intraosseous (IO) fluids should be placed wide open with pressure bags.
- C. If a palpable pulse becomes present:
 1. Re-assess for and control external hemorrhage.
 2. Administer TXA as indicated per PD# 8065 – Hemorrhage Control.
 3. Titrate normal saline to SBP $\geq$ 90 mmHg or palpable peripheral pulses.

Cross Reference:

PD# 2033 – Determination of Death
PD# 2085 – Do Not Resuscitate
PD# 8015 – Trauma
PD# 8020 – Respiratory Distress - Airway Management
PD# 8024 – Cardiac Dysrhythmias
PD# 8026 – Respiratory Distress
PD# 8031 – Non-Traumatic Cardiac Arrest
PD# 8044 – Spinal Motion Restrictions
PD# 8065 – Hemorrhage Control

